

Big Ramsey degrees of 3-uniform hypergraphs are finite

David Chodounský

Charles University

Joint work with M. Balko, J. Hubička, M. Konečný, and L. Vena



Martin Balko, David Chodounský, Jan Hubička,
Matěj Konečný, Lluís Vena,
Big Ramsey degrees of 3-uniform hypergraphs are finite,
<https://arxiv.org/abs/2008.00268>

Definition

Let \mathbf{A} be a countable structure. We say that \mathbf{A} has *finite big Ramsey degrees* if for every $n \in \omega$ there is $D(n) \in \omega$ such that for every finite coloring of $[\mathbf{A}]^n$ there is a copy \mathbf{B} of \mathbf{A} (inside of \mathbf{A}) such that $[\mathbf{B}]^n$ has at most $D(n)$ colors.

Example

- ▶ $(\omega, \text{no structure})$ (Ramsey)
- ▶ $(\mathbb{Q}, <)$ (Laver, Devlin)
- ▶ Random (Rado) graph (Todorčević, Sauer)
- ▶ Triangle free Henson graph \mathbb{H}_3 (Dobrinen, see also Hubička)
- ▶ Random 3-hypergraph (BHChKV)

Definition

A structure $\mathbf{A} \in \mathcal{C}$ is universal (for a class of structures \mathcal{C}) if \mathbf{A} contains a copy of every $\mathbf{B} \in \mathcal{C}$.

Definition

A structure $\mathbf{A} \in \mathcal{C}$ is universal (for a class of structures \mathcal{C}) if \mathbf{A} contains a copy of every $\mathbf{B} \in \mathcal{C}$.

Proposition

If $\mathbf{A}, \mathbf{B} \in \mathcal{C}$ are both universal for \mathcal{C} and \mathbf{A} has finite big Ramsey degrees, then \mathbf{B} also has finite big Ramsey degrees.

Trees

- ▶ rooted
- ▶ height at most $\omega \dots h(T) \leq \omega$
- ▶ finitely branching
- ▶ balanced (no short branches)
- ▶ n -th level of $T \dots T(n)$
- ▶ initial subtree $\dots T(<n)$
- ▶ set of immediate successors of s in $T \dots isu_T(s)$

Definition

A subtree S of T of height $h(S) = n \in \omega + 1$ is a *strong subtree* if

- ▶ $\forall n < h(S) \exists m < h(T)$ such that $S(n) \subseteq T(m)$,
- ▶ $\forall s \in S \forall t \in isu_T(s) \exists!(s' \in S, s' \geq t, s' \in isu_S(s))$,
unless $isu_S(s) = \emptyset$.

We write $S \in STR_n(T)$.

Definition

A subtree S of T of height $h(S) = n \in \omega + 1$ is a *strong subtree* if

- ▶ $\forall n < h(S) \exists m < h(T)$ such that $S(n) \subseteq T(m)$,
- ▶ $\forall s \in S \forall t \in isu_T(s) \exists!(s' \in S, s' > t, s' \in isu_S(s))$,
unless $isu_S(s) = \emptyset$.

We write $S \in STR_n(T)$.

Definition

A subtree S of T of height $h(S) = n \in \omega + 1$ is a *strong subtree* if

- ▶ $\forall n < h(S) \exists m < h(T)$ such that $S(n) \subseteq T(m)$,
- ▶ $\forall s \in S \forall t \in \text{isu}_T(s) \exists!(s' \in S, s' > t, s' \in \text{isu}_S(s))$,
unless $\text{isu}_S(s) = \emptyset$.

We write $S \in \text{STR}_n(T)$.

If $S \in \text{STR}_n(T)$ and $R \in \text{STR}_m(S)$, then $R \in \text{STR}_m(T)$.

Theorem (Milliken, simple version)

If T is a tree of height ω , $n \in \omega$, and $\chi: \text{STR}_n(T) \rightarrow k$ is a finite coloring, then there exists $S \in \text{STR}_\omega(T)$ such that χ is monochromatic on $\text{STR}_n(S)$.

Trees, examples

Example

$\mathbf{T}_B = 2^{<\omega}$, the binary tree

Observation

If $S \in STR_n(\mathbf{T}_B)$, then S is isomorphic to $\mathbf{T}_B(<n)$.

Example

$\mathbf{T}_M = \bigcup \{ 2^{n \times n} : n \in \omega \}$, ordered by extension. The tree of matrices.

$\mathbf{T}_{SM} \subset \mathbf{T}_M$, the tree of sub-diagonal matrices.

If $A \in \mathbf{T}_{SM}$ and $A(i, j) \neq 0$, then $i < j$.

For $A \in \mathbf{T}_M(n)$ we write $|A| = n$.

Random graph has finite big Ramsey degrees

For $s, t \in \mathbf{T}_B$ define $E(s, t)$ if $|s| < |t|$ and $t(|s|) = 1$.

Proposition

The graph (\mathbf{T}_B, E) is universal (for the class of all countable graphs).

Random graph has finite big Ramsey degrees

For $s, t \in \mathbf{T}_B$ define $E(s, t)$ if $|s| < |t|$ and $t(|s|) = 1$.

Proposition

The graph (\mathbf{T}_B, E) is universal (for the class of all countable graphs).

Observation

If $S \in STR_\omega(\mathbf{T}_B)$, then (S, E) is a copy of (\mathbf{T}_B, E)
(both as a graph and as a tree).

Random graph has finite big Ramsey degrees

For $s, t \in \mathbf{T}_B$ define $E(s, t)$ if $|s| < |t|$ and $t(|s|) = 1$.

Proposition

The graph (\mathbf{T}_B, E) is universal (for the class of all countable graphs).

Observation

If $S \in STR_\omega(\mathbf{T}_B)$, then (S, E) is a copy of (\mathbf{T}_B, E)
(both as a graph and as a tree).

Lemma

*For every $n \in \omega$ and $a \in [\mathbf{T}_B]^n$ there exists
 $S_B \in STR_{2n}(\mathbf{T}_B)$ such that $a \subset S_B$.*

Random graph has finite big Ramsey degrees

For $s, t \in \mathbf{T}_B$ define $E(s, t)$ if $|s| < |t|$ and $t(|s|) = 1$.

Proposition

The graph (\mathbf{T}_B, E) is universal (for the class of all countable graphs).

Observation

If $S \in STR_\omega(\mathbf{T}_B)$, then (S, E) is a copy of (\mathbf{T}_B, E)
(both as a graph and as a tree).

Lemma

*For every $n \in \omega$ and $a \in [\mathbf{T}_B]^n$ there exists
 $S_B \in STR_{2n}(\mathbf{T}_B)$ such that $a \subset S_B$.*

Proof

Given finite coloring $\chi: [\mathbf{T}_B]^n \rightarrow k$.

Induces finite coloring $\bar{\chi}: STR_{2n}(\mathbf{T}_B) \rightarrow k^{(2^{2n})^n}$.

Use Milliken's theorem.

Universal 3-hypergraphs have finite big Ramsey degrees

For $A, B, C \in \mathbf{T}_{SM}$ define $E(A, B, C)$ is $|A| < |B| < |C|$ and $C(|A|, |B|) = 1$.

Universal 3-hypergraphs have finite big Ramsey degrees

For $A, B, C \in \mathbf{T}_{SM}$ define $E(A, B, C)$ is $|A| < |B| < |C|$ and $C(|A|, |B|) = 1$.

Proposition

The hypergraph (\mathbf{T}_{SM}, E) is universal (for countable 3-hypergraphs).

Observation

If $S \in STR_\omega(\mathbf{T}_{SM})$, then (S, E) is **not** a copy of (\mathbf{T}_{SM}, E) .
(But we can find a copy of \mathbf{T}_{SM} inside S .)

Product trees

$\mathbf{T}_{SM} \otimes \mathbf{T}_B$... the product tree

Definition

We say that $S_{SM} \otimes S_B \in STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$

($S_{SM} \otimes S_B$ is a strong subtree of $\mathbf{T}_{SM} \otimes \mathbf{T}_B$) if

- ▶ $S_{SM} \in STR_k(\mathbf{T}_{SM})$,
- ▶ $S_B \in STR_k(\mathbf{T}_B)$, and
- ▶ $\forall n \in k \exists m \in \omega$ such that $S_{SM}(n) \subseteq \mathbf{T}_{SM}(m)$ and $S_B(n) \subseteq \mathbf{T}_B(m)$.

Product trees

$\mathbf{T}_{SM} \otimes \mathbf{T}_B$... the product tree

Definition

We say that $S_{SM} \otimes S_B \in STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$
($S_{SM} \otimes S_B$ is a strong subtree of $\mathbf{T}_{SM} \otimes \mathbf{T}_B$) if

- ▶ $S_{SM} \in STR_k(\mathbf{T}_{SM})$,
- ▶ $S_B \in STR_k(\mathbf{T}_B)$, and
- ▶ $\forall n \in k \exists m \in \omega$ such that $S_{SM}(n) \subseteq \mathbf{T}_{SM}(m)$ and $S_B(n) \subseteq \mathbf{T}_B(m)$.

Theorem (Milliken, special case)

If $k \in \omega$ and $\chi: STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B) \rightarrow k$ is a finite coloring, then there exists $S_{SM} \otimes S_B \in STR_\omega(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$ such that χ is monochromatic on $STR_k(S_{SM} \otimes S_B)$.

Valuations

Suppose $S_{SM} \otimes S_B \in STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$ for some $k \in \omega + 1$.

We define the tree $val(S_{SM} \times S_B) \subseteq S_{SM}$ by induction:

- ▶ The root of $val(S_{SM} \times S_B)$ is the root of S_{SM} .
- ▶ If $A \in val(S_{SM} \otimes S_B)$, $t \in S_B(|A|)$, $C \in isu_{S_{SM}}(A)$, and $C > A \hat{\ } t$, then $A \in val(S_{SM} \otimes S_B)$.

Valuations

Suppose $S_{SM} \otimes S_B \in STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$ for some $k \in \omega + 1$.

We define the tree $val(S_{SM} \times S_B) \subseteq S_{SM}$ by induction:

- ▶ The root of $val(S_{SM} \times S_B)$ is the root of S_{SM} .
- ▶ If $A \in val(S_{SM} \otimes S_B)$, $t \in S_B(|A|)$, $C \in isu_{S_{SM}}(A)$, and $C > A \hat{\ } t$, then $A \in val(S_{SM} \otimes S_B)$.

Observation

If $S_{SM} \otimes S_B \in STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$,

then $(val(S_{SM} \otimes S_B), E)$ is a copy of $(\mathbf{T}_{SM}(< k), E)$

(both as a hypergraph and as a tree).

Lemma (false but fixable)

For every $n \in \omega$ there is $f(n) \in \omega$

such that for every $a \in [\mathbf{T}_{SM}]^n$ there exists

$S_{SM} \otimes S_B \in STR_{f(n)}(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$ such that $a \subset val(S_{SM} \otimes S_B)$.

Observation

If $S_{SM} \otimes S_B \in STR_k(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$,
then $(val(S_{SM}, S_B), E)$ is a copy of $(\mathbf{T}_{SM}(< k), E)$
(both as a hypergraph and as a tree).

Lemma (false but fixable)

For every $n \in \omega$ there is $f(n) \in \omega$
such that for every $a \in [\mathbf{T}_{SM}]^n$ there exists
 $S_{SM} \otimes S_B \in STR_{f(n)}(\mathbf{T}_{SM} \otimes \mathbf{T}_B)$ such that $a \subset val(S_{SM}, S_B)$.

Proof

Given finite coloring $\chi: [\mathbf{T}_{SM}]^n \rightarrow k$.

Induces finite coloring $\bar{\chi}: STR_{f(n)}(\mathbf{T}_{SM} \otimes \mathbf{T}_B) \rightarrow K$

(look at colors on valuations).

Use Milliken's theorem.